

DENSE SUBSETS OF BOUNDARIES OF CAT(0) GROUPS

TETSUYA HOSAKA

ABSTRACT. In this paper, we study dense subsets of boundaries of CAT(0) groups. Suppose that a group G acts geometrically on a CAT(0) space X and suppose that there exists an element $g_0 \in G$ such that (1) Z_{g_0} is finite, (2) $X \setminus F_{g_0}$ is not connected, and (3) each component of $X \setminus F_{g_0}$ is convex and not g_0 -invariant, where Z_{g_0} is the centralizer of g_0 and F_{g_0} is the fixed-point set of g_0 in X (that is, $Z_{g_0} = \{h \in G \mid g_0 h = h g_0\}$ and $F_{g_0} = \{x \in X \mid g_0 x = x\}$). Then we show that each orbit $G\alpha$ is dense in the boundary ∂X (i.e. ∂X is minimal) and the set $\{g^\infty \mid g \in G, o(g) = \infty\}$ is also dense in the boundary ∂X . We obtain an application for dense subsets on the boundary of a Coxeter system.

1. INTRODUCTION AND PRELIMINARIES

In this paper, we study dense subsets of boundaries of CAT(0) groups. Definitions and basic properties of CAT(0) spaces and their boundaries are found in [1]. A *geometric* action on a CAT(0) space is an action by isometries which is proper ([1, p.131]) and cocompact. We note that every CAT(0) space on which some group acts geometrically is a proper space ([1, p.132]).

Suppose that a group G acts on a compact metric space Y by homeomorphisms. Then Y is said to be *minimal*, if every orbit Gy is dense in Y .

For a negatively curved group G and the boundary ∂G of G , by an easy argument, we can show that $G\alpha$ is dense in ∂G for each $\alpha \in \partial G$, that is, ∂G is minimal.

Suppose that a group G acts geometrically on a CAT(0) space X . Then G acts on the boundary ∂X by homeomorphisms. The boundary ∂X need not be minimal in general. Indeed, there exist some examples of Coxeter systems whose boundaries are not minimal (cf. [3] and [5]).

Date: September 19, 2005.

2000 Mathematics Subject Classification. 57M07.

Key words and phrases. boundaries of CAT(0) groups.

Partly supported by the Grant-in-Aid for Young Scientists (B), The Ministry of Education, Culture, Sports, Science and Technology, Japan. (No. 15740029).

In [3] and [5], we have investigated dense subsets (and dense orbits) of the boundary of a Coxeter system. In this paper, we study dense subsets of boundaries of CAT(0) groups. The purpose of this paper is to prove the following theorem.

Theorem 1.1. *Suppose that a group G acts geometrically on a CAT(0) space X . If there exists an element $g_0 \in G$ such that*

- (1) Z_{g_0} is finite,
- (2) $X \setminus F_{g_0}$ is not connected, and
- (3) each component of $X \setminus F_{g_0}$ is convex and not g_0 -invariant,

then each orbit $G\alpha$ is dense in the boundary ∂X (i.e. ∂X is minimal) and the set $\{g^\infty \mid g \in G, o(g) = \infty\}$ is also dense in ∂X . Here Z_{g_0} is the centralizer of g_0 and F_{g_0} is the fixed-point set of g_0 in X .

We give the proof of this theorem in Section 2, and we give some applications of this theorem and introduce some open problems about dense orbits of boundaries of CAT(0) groups in Section 3.

2. PROOF OF THE MAIN THEOREM

We first show some results about CAT(0) groups and their boundaries needed in the proof of Theorem 1.1.

Suppose that a group G acts geometrically on a CAT(0) space X . For an element $g \in G$, we define Z_g as the centralizer of g and define F_g and $\mathcal{F}_g$ as the fixed-point sets of g in X and ∂X respectively, that is,

$$\begin{aligned} Z_g &= \{h \in G \mid gh = hg\}, \\ F_g &= \{x \in X \mid gx = x\} \text{ and} \\ \mathcal{F}_g &= \{\alpha \in \partial X \mid g\alpha = \alpha\}. \end{aligned}$$

Also for a subset $A \subset G$, the limit set $L(A)$ of A is defined as

$$L(A) = \overline{Ax_0} \cap \partial X,$$

where $x_0 \in X$ and $\overline{Ax_0}$ is the closure of the orbit Ax_0 in $X \cup \partial X$. We note that the limit set $L(A)$ is determined by A and not depend on the point x_0 .

The following theorem has been proved in [6].

Theorem 2.1. *Suppose that a group G acts geometrically on a CAT(0) space X . For $g \in G$, the equality $\mathcal{F}_g = L(Z_g)$ holds. Moreover, $\mathcal{F}_g = \emptyset$ if and only if Z_g is finite.*

Here if F_g is unbounded then $\mathcal{F}_g \neq \emptyset$. Hence if Z_g is finite then F_g is bounded.

We show the following basic lemma.

Lemma 2.2. *Suppose that a group G acts geometrically on a $CAT(0)$ space X . Let A be a subset of the boundary ∂X , let $x_0 \in X$ and let $M, N > 0$. If for each $g \in G$ such that $d(x_0, gx_0) > M$, there exists $\alpha \in A$ such that $d(gx_0, \text{Im } \xi_\alpha) \leq N$, then A is dense in ∂X . Here ξ_α is the geodesic ray in X such that $\xi_\alpha(0) = x_0$ and $\xi_\alpha(\infty) = \alpha$.*

Proof. Suppose that for each $g \in G$ such that $d(x_0, gx_0) > M$, there exists $\alpha \in A$ such that $d(gx_0, \text{Im } \xi_\alpha) \leq N$.

Let $\beta \in \partial X$ and let ξ_β be the geodesic ray in X such that $\xi_\beta(0) = x_0$ and $\xi_\beta(\infty) = \beta$. To prove that A is dense in ∂X , we show that for each $\epsilon > 0$ and $R > 0$, there exists $\alpha \in A$ such that $d(\xi_\beta(R), \text{Im } \xi_\alpha) < \epsilon$.

Let $\epsilon > 0$ and $R > 0$. Since the action of G on X is cocompact, $B(Gx_0, N') = X$ for some $N' > 0$. Let R' be a number such that

$$R' > \max \left\{ \frac{R(N + N')}{\epsilon}, M + N' \right\}.$$

Since $\xi_\beta(R') \in X = B(Gx_0, N')$, there exists $g \in G$ such that $d(\xi_\beta(R'), gx_0) \leq N'$. Then

$$\begin{aligned} d(x_0, gx_0) &\geq d(x_0, \xi_\beta(R')) - d(\xi_\beta(R'), gx_0) \\ &\geq R' - N' > (M + N') - N' = M. \end{aligned}$$

Hence $d(x_0, gx_0) > M$, and there exists $\alpha \in A$ such that $d(gx_0, \text{Im } \xi_\alpha) \leq N$. Then

$$d(\xi_\beta(R'), \text{Im } \xi_\alpha) \leq d(\xi_\beta(R'), gx_0) + d(gx_0, \text{Im } \xi_\alpha) \leq N' + N.$$

Hence

$$\begin{aligned} d(\xi_\beta(R), \text{Im } \xi_\alpha) &\leq \frac{R}{R'} d(\xi_\beta(R'), \text{Im } \xi_\alpha) \\ &\leq \frac{R}{R'} (N + N') < \epsilon. \end{aligned}$$

Thus A is dense in ∂X . □

The following lemma is known.

Lemma 2.3 (cf. [1, Proposition II.6.2(2)]). *Let X be a $CAT(0)$ space and let g and h be isometries of X . Then $gF_h = F_{ghg^{-1}}$, where F_h is the fixed-point set of h in X .*

Using the above results, we prove Theorem 1.1.

Proof of Theorem 1.1. Suppose that a group G acts geometrically on a $CAT(0)$ space X and suppose that there exists an element $g_0 \in G$ such that

- (1) Z_{g_0} is finite,
- (2) $X \setminus F_{g_0}$ is not connected, and
- (3) each component of $X \setminus F_{g_0}$ is convex and not g_0 -invariant.

Then F_{g_0} is bounded by (1) and Theorem 2.1. Let x_0 be a basepoint of X and let N be a number such that $F_{g_0} \subset B(x_0, N)$.

To prove that every orbit $G\alpha$ is dense in ∂X , by Lemma 2.2, we show that for each $\alpha \in \partial X$ and $g \in G$, there exists $h \in G$ such that $d(gx_0, \text{Im } \xi_{h\alpha}) \leq N$, where $\xi_{h\alpha}$ is the geodesic ray in X such that $\xi_{h\alpha}(0) = x_0$ and $\xi_{h\alpha}(\infty) = h\alpha$.

Let $\alpha \in \partial X$ and $g \in G$.

If $\text{Im } \xi_\alpha \cap gF_{g_0} \neq \emptyset$, then

$$d(gx_0, \text{Im } \xi_\alpha) \leq d(gx_0, \text{Im } \xi_\alpha \cap gF_{g_0}) \leq N,$$

because $gF_{g_0} \subset B(gx_0, N)$.

Suppose that $\text{Im } \xi_\alpha \cap gF_{g_0} = \emptyset$. Let gX' be the component of $X \setminus gF_{g_0}$ such that $x_0 \in gX'$. Since $\text{Im } \xi_\alpha \cap gF_{g_0} = \emptyset$, we have that $\text{Im } \xi_\alpha \subset gX'$. Then $gg_0g^{-1}(\text{Im } \xi_\alpha) \subset gg_0g^{-1}(gX') = gg_0X'$. Here $gX' \neq gg_0X'$ because $X' \neq g_0X'$ by non- g_0 -invariantness of X' which is a component of $X \setminus F_{g_0}$. Hence $gg_0g^{-1}\alpha \in \partial(gg_0X')$. On the other hand, $x_0 \in gX'$. Thus $\text{Im } \xi_{gg_0g^{-1}\alpha} \cap gF_{g_0} \neq \emptyset$, where $\xi_{gg_0g^{-1}\alpha}$ is the geodesic ray in X such that $\xi_{gg_0g^{-1}\alpha}(0) = x_0$ and $\xi_{gg_0g^{-1}\alpha}(\infty) = gg_0g^{-1}\alpha$. Hence

$$d(gx_0, \text{Im } \xi_{gg_0g^{-1}\alpha}) \leq d(gx_0, \text{Im } \xi_{gg_0g^{-1}\alpha} \cap gF_{g_0}) \leq N,$$

since $gF_{g_0} \subset B(gx_0, N)$.

Therefore every orbit $G\alpha$ is dense in ∂X .

Next we prove that the set $\{h^\infty \mid h \in G, o(h) = \infty\}$ is dense in ∂X . By Lemma 2.2, we show that for each $g \in G$ such that $d(x_0, gx_0) > 2N$, there exists $h \in G$ such that $o(h) = \infty$ and $d(gx_0, \text{Im } \xi_{h^\infty}) \leq N$.

Let $g \in G$ such that $d(x_0, gx_0) > 2N$.

We first suppose that $[x_0, gx_0] \cap gF_{g_0} = \emptyset$, where $[x_0, gx_0]$ is the geodesic from x_0 to gx_0 in X . Then

$$\begin{aligned} d(gF_{g_0}, gg_0gF_{g_0}) &= d(F_{g_0}, g_0gF_{g_0}) \\ &= d(g_0^{-1}F_{g_0}, gF_{g_0}) \\ &= d(F_{g_0}, gF_{g_0}) \\ &\geq d(x_0, gx_0) - 2N \\ &> 0, \end{aligned}$$

because $g_0F_{g_0} = F_{g_0}$, $F_{g_0} \subset B(x_0, N)$ and $d(x_0, gx_0) > 2N$. Hence $gF_{g_0} \cap gg_0gF_{g_0} = \emptyset$. Thus $(gg_0)^i[x_0, gx_0] \cap (gg_0)^igF_{g_0} = \emptyset$ and

$(gg_0)^i gF_{g_0} \cap (gg_0)^{i+1} gF_{g_0} = \emptyset$ for each $i \in \{0, 1, 2, \dots\}$. We consider

$$Y = X \setminus \bigcup \{(gg_0)^i gF_{g_0} \mid i = 0, 1, 2, \dots\}.$$

Let Y_i be the component of Y such that $(gg_0)^i x_0 \in Y_i$ for each $i \in \{0, 1, 2, \dots\}$. Then $Y_i \neq Y_j$ if $i \neq j$ by the above argument. Hence $o(gg_0) = \infty$. We consider $(gg_0)^\infty$ and the geodesic ray $\xi_{(gg_0)^\infty}$. Since the sequence $\{(gg_0)^i\}$ converges to $(gg_0)^\infty$ and $(gg_0)^i x_0 \in Y_i$ for each i , we obtain that $\text{Im } \xi_{(gg_0)^\infty} \cap (gg_0)^i gF_{g_0} \neq \emptyset$ for each i . In particular, $\text{Im } \xi_{(gg_0)^\infty} \cap gF_{g_0} \neq \emptyset$. Thus

$$d(gx_0, \text{Im } \xi_{(gg_0)^\infty}) \leq d(gx_0, \text{Im } \xi_{(gg_0)^\infty} \cap gF_{g_0}) \leq N,$$

since $gF_{g_0} \subset B(gx_0, N)$.

Next we suppose that $[x_0, gx_0] \cap gF_{g_0} \neq \emptyset$. Let gX' be the component of $X \setminus gF_{g_0}$ such that $x_0 \in gX'$. Since X' is not g_0 -invariant, there exists a number k such that $gg_0^k x_0 \in gX'$. Then $[x_0, gg_0^k x_0] \cap gF_{g_0} = \emptyset$ because $x_0, gg_0^k x_0 \in gX'$ and gX' is convex. Here

$$\begin{aligned} d(gg_0^k F_{g_0}, (gg_0^k)g_0(gg_0^k)F_{g_0}) &= d(F_{g_0}, g_0(gg_0^k)F_{g_0}) \\ &= d(F_{g_0}, g_0 gF_{g_0}) \\ &= d(g_0^{-1}F_{g_0}, gF_{g_0}) \\ &= d(F_{g_0}, gF_{g_0}) \\ &\geq d(x_0, gx_0) - 2N \\ &> 0. \end{aligned}$$

Hence $gg_0^k F_{g_0} \cap (gg_0^k)g_0(gg_0^k)F_{g_0} = \emptyset$. By the same argument as the above one, we have that $\text{Im } \xi_{((gg_0^k)g_0)^\infty} \cap gg_0^k F_{g_0} \neq \emptyset$. Since $g_0^k F_{g_0} = F_{g_0}$, $\text{Im } \xi_{(gg_0^{k+1})^\infty} \cap gF_{g_0} \neq \emptyset$. Thus

$$d(gx_0, \text{Im } \xi_{(gg_0^{k+1})^\infty}) \leq d(gx_0, \text{Im } \xi_{(gg_0^{k+1})^\infty} \cap gF_{g_0}) \leq N.$$

Therefore the set $\{h^\infty \mid h \in G, o(h) = \infty\}$ is dense in ∂X . $\square$

3. APPLICATIONS AND PROBLEMS

In [4], we have defined a *reflection* of a geodesic space. We say that an isometry r of a geodesic space X is a *reflection* of X , if

- (1) r^2 is the identity of X ,
- (2) $X \setminus F_r$ has exactly two convex connected components X_r^+ and X_r^- ,
- (3) $rX_r^+ = X_r^-$ and
- (4) $\text{Int } F_r = \emptyset$,

where F_r is the fixed-point set of r .

We obtain the following corollary from Theorem 1.1.

Corollary 3.1. *Suppose that a group G acts geometrically on a $CAT(0)$ space X . If there exists a reflection $r \in G$ of X such that Z_r is finite, then each orbit $G\alpha$ is dense in the boundary ∂X (i.e. ∂X is minimal) and the set $\{g^\infty \mid g \in G, o(g) = \infty\}$ is dense in ∂X . Here Z_r is the centralizer of r .*

For a Coxeter system (W, S) and the Davis complex $\Sigma(W, S)$ which is a $CAT(0)$ space ([2] and [7]), each $s \in S$ is a reflection of $\Sigma(W, S)$. Hence we obtain the following corollary.

Corollary 3.2. *Let (W, S) be a Coxeter system and let $\Sigma(W, S)$ be the Davis complex of (W, S) . If there exists $s \in S$ such that Z_s is finite, then each orbit $W\alpha$ is dense in the boundary $\partial\Sigma(W, S)$ (i.e. $\partial\Sigma(W, S)$ is minimal) and the set $\{w^\infty \mid w \in W, o(w) = \infty\}$ is dense in $\partial\Sigma(W, S)$.*

The following problems are open.

Problem. Does there exist a Coxeter system (W, S) such that some orbit $W\alpha$ is dense in $\partial\Sigma(W, S)$ and $\partial\Sigma(W, S)$ is not minimal?

Problem. Suppose that a group G acts geometrically on two $CAT(0)$ spaces X and X' . Is it the case that ∂X is minimal if and only if $\partial X'$ is minimal?

Problem. Suppose that a group G acts geometrically on a $CAT(0)$ space X . Is it always the case that the set $\{g^\infty \mid g \in G, o(g) = \infty\}$ is dense in ∂X ?

REFERENCES

- [1] M. R. Bridson and A. Haefliger, *Metric spaces of non-positive curvature*, Springer-Verlag, Berlin, 1999.
- [2] M. W. Davis, *Nonpositive curvature and reflection groups*, in Handbook of geometric topology (Edited by R. J. Daverman and R. B. Sher), pp. 373–422, North-Holland, Amsterdam, 2002.
- [3] T. Hosaka, *Dense subsets of the boundary of a Coxeter system*, Proc. Amer. Math. Soc. **132** (2004), 3441–3448.
- [4] T. Hosaka, *Reflection groups of geodesic spaces and Coxeter groups*, Topology Appl., to appear.
- [5] T. Hosaka, *On dense orbits in the boundary of a Coxeter system*, J. Kyoto Univ., to appear.
- [6] T. Hosaka, *On fixed-point sets in the boundary of a $CAT(0)$ space*, preprint.
- [7] G. Moussong, *Hyperbolic Coxeter groups*, Ph.D. thesis, The Ohio State University, 1988.

DEPARTMENT OF MATHEMATICS, UTSUNOMIYA UNIVERSITY, UTSUNOMIYA, 321-8505, JAPAN

E-mail address: hosaka@cc.utsunomiya-u.ac.jp